arXiv:astro-ph/0402363v1 16 Feb 2004

Spectral Statistics and Luminosity Function of a Hard X-ray Complete Sample of Brightest AGNs

K. Shinozaki[1],*) T. Miyaji[2], Y. Ishisaki[1], Y. Ueda[3], Y. Ogasaka[4],
K. Hayashida[5], H. Awaki[6]

[1] *Experimental Astrophysics Group, Department of Physics, Tokyo Metropolitan University Tokyo, Japan*
[2] *Department of Physics, Carnegie Mellon University*
[3] *ISAS/JAXA*
[4] *Department of Astrophysics, Nagoya University*
[5] *Department of Astrophysics, Osaka University*
[6] *Department of Physics, Ehime University*

Investigating the X-ray luminosity function of AGNs in different redshifts is very important in understanding growth of super massive black holes in centers of galaxies over the cosmological time scale. As a part of this, we are investigating the statistics of the X-ray spectral properties of a complete flux-limited sample of bright AGNs. The purpose of this investigation is to provide the present-day constraint on the evolution of absorbed and unabsorbed AGNs, and to constrain the AGN population synthesis model of the total X-ray background.



Fig. 1. Spatial ditribution of AGNs in our two samples. Circles denote the Sample-1 (Piccinotti) and Squares denote the Sample-2 (Grossan). Dashed lines correspond to b=±20°, and Dotted region represents the selected resion for the sample-2.



Fig. 2. Hard X-ray luminosity function of all AGNs (Square), X-ray unabsorbed AGNs (Circle), and absorbed AGNs (Triangle).

We defined 28 AGNs in Picinotti et al. (1982),[1] as a Sample-1, selecting the R15 count rate > 1.25 c s^{-1} $\sim 3 \times 10^{-11}$ erg cm^{-2} s^{-1}. The Galactic latitude $|\mathrm{b}| < 20°$ and the LMC region were excluded. Two sources which are also in Sample-2 were excluded. This sample covers 65.5 % of the sky. We also defined Sample-2, which contains 20 sources from Grossan (1992).[2] This sample was selected from the MC-

---

*) shino@phys.metro-u.ac.jp

typeset using $\mathcal{PTP}$TEX.cls ⟨Ver.0.89⟩

LASS catalog of HEAO-1 A1/A3 sources and with a flux-limit of $\sim 1.7 \times 10^{-11}$ erg $\mathrm{cm}^{-2}\ \mathrm{s}^{-1}$, inside the 55°-radius from the NEP. As a result, we selected 48 sources as a complete flux-limited sample with $z < 0.4$ (Fig. 1). Spectral studies have been made using ASCA (14 sources) or XMM-Newton (22 sources) observations by ourselves, and other 12 sources were derived from literature[3)−.11)] These two samples are also used to calculate a wide range HXLF in Ueda et al. (2003).[13)]

The spectral measurements enabled us to construct local HXLF for absorbed ($\mathrm{Log}\, N_{\mathrm{H}}[\mathrm{cm}^{-2}] > 21.5$) and unabsorbed AGNs separately. We calculated each HXLF with the conventional $\Sigma V_{\mathrm{a}}^{-1}$ method[12)] (Fig. 2). We found a high luminosity cut-off for the HXLF of the absorbed AGNs compared with those of unabsorbed AGNs and all AGNs. In the $L_{\mathrm{x}} - N_{\mathrm{H}}$ plot (Fig. 3), there are no AGNs in the region where $\mathrm{Log}\, L_{\mathrm{X}}[\mathrm{erg/s}] > 44.5$ and $\mathrm{Log}\, N_{\mathrm{H}}[\mathrm{cm}^{-2}] > 21.5$. We would have 9-10 AGNs in this region if the shapes of the absorbed and unabsorbed AGN HXLF were identical. We found that the observed flux with ASCA or XMM-Newton was smaller than that with HEAO-1 by a factor of 0.42 on average, which was expected for re-observation of sources with a variability-amplitude of a factor ~2.6 (Fig. 4).



Fig. 3. Relation between hard X-ray luminosity and intrinsic N_{H}.



Fig. 4. Comparison of observed flux between ASCA or XMM-Newton, and HEAO-1.

References

1) Piccinotti, G., Mushotzky, R.F., Boldt, E.A., et al. Astrophys. J. **1982** (253), 485
2) Grossan B.A. 1992, PhD Thesis,MIT
3) Reynolds, C.S. Royal Astronomical Society, Monthly Notices, 1997, 286, 513
4) Turner, T.J., George, I.M., Nandra, K., and Mushotzky, R.F. Astrophys. J. Supplement, 1997, 113, 23
5) Guainazzi, M. Astronomy and Astrophys, 2003, 401, 903
6) George, I.M., Turner, T.J., Netzer, H., et al. Astrophys. J. Supplement, 1998, 114, 73
7) Blustin, A.J., Branduardi-Raymont, G., Behar, E., et al. Astronomy and Astrophys, 2002, 392, 453
8) Gondoin, P., Orr, A., and Lumb, D. Astronomy and Astrophys, 2003, 398, 967
9) Gondoin, P., Orr, A., Lumb, D., and Siddiqui, H. Astronomy and Astrophys, 2003, 397, 883
10) Malizia, A., Malaguti, G., Bassani, L., et al. Astronomy and Astrophys, 2002, 394, 801
11) Turner, T.J., Netzer, H., and George, I.M. Astrophys. J. **1996** (463), 134
12) Miyaji, T., Hasinger, G., and Schmidt, M. Astronomy and Astrophys 2001, 369, 49a
13) Ueda, Y., Akiyama, M., Ohta, K., and Miyaji, T. Astrophys. J. **2003** (598), 886